





















Erythema multiforme-like pattern of lesions in a patient with pemphigoid gestationis























































Erythema multiforme is a skin condition characterized by red, target-like lesions. It is often caused by a reaction to certain medications or infections. The lesions typically appear on the extremities and can be accompanied by itching or pain. The condition is usually self-limiting and resolves within a few weeks. Treatment is primarily symptomatic, focusing on relieving discomfort and avoiding the triggering agent. In severe cases, systemic corticosteroids may be used. The condition is more common in children and young adults.

erythema multiforme is a skin condition characterized by red, target-like lesions. It is often caused by a reaction to certain medications or infections. The lesions typically appear on the extremities and can be accompanied by itching or pain. The condition is usually self-limiting and resolves within a few weeks. Treatment is primarily symptomatic, focusing on relieving discomfort and avoiding the triggering agent. In severe cases, systemic corticosteroids may be used. The condition is more common in children and young adults.

erythema multiforme is a skin condition characterized by red, target-like lesions. It is often caused by a reaction to certain medications or infections. The lesions typically appear on the extremities and can be accompanied by itching or pain. The condition is usually self-limiting and resolves within a few weeks. Treatment is primarily symptomatic, focusing on relieving discomfort and avoiding the triggering agent. In severe cases, systemic corticosteroids may be used. The condition is more common in children and young adults.