











Acrokeratosis of the face and hands is a paraneoplastic syndrome that is associated with squamous cell carcinoma of the lung. It is characterized by the presence of thick, yellowish, scaly crusts on the fingers and toes, and on the face, particularly around the nose and mouth. The condition is usually self-limiting and resolves after the primary tumor is treated. The pathogenesis is thought to be related to the production of autoantibodies against the keratinocytes of the skin. The condition is also associated with other paraneoplastic syndromes, such as hypercalcemia, weight loss, and fatigue. The diagnosis is based on the clinical presentation and the presence of the characteristic skin lesions. The treatment is primarily supportive and focuses on the removal of the crusts and the use of topical steroids to reduce inflammation. The prognosis is generally good, as the condition usually resolves after the primary tumor is treated. However, the condition can be a sign of a more serious underlying malignancy, so it is important to investigate the cause of the skin lesions. The condition is also associated with other paraneoplastic syndromes, such as hypercalcemia, weight loss, and fatigue. The diagnosis is based on the clinical presentation and the presence of the characteristic skin lesions. The treatment is primarily supportive and focuses on the removal of the crusts and the use of topical steroids to reduce inflammation. The prognosis is generally good, as the condition usually resolves after the primary tumor is treated. However, the condition can be a sign of a more serious underlying malignancy, so it is important to investigate the cause of the skin lesions.